



LARIAM

brand of

mefloquine hydrochloride

TABLETS


R_x only

DESCRIPTION

Lariam (mefloquine hydrochloride) is an antimalarial agent available as 250-mg tablets of mefloquine hydrochloride (equivalent to 228.0 mg of the free base) for oral administration.

Mefloquine hydrochloride is a 4-quinolinemethanol derivative with the specific chemical name of (R*, S*)-(±)-α-2-piperidiny-2,8-bis(trifluoromethyl)-4-quinolinemethanol hydrochloride. It is a 2-aryl substituted chemical structural analog of quinine. The drug is a white to almost white crystalline compound, slightly soluble in water.

Mefloquine hydrochloride has a calculated molecular weight of 414.78 and the following structural formula:



The inactive ingredients are ammonium-calcium alginate, corn starch, crospovidone, lactose, magnesium stearate, microcrystalline cellulose, poloxamer #331, and talc.

CLINICAL PHARMACOLOGY

Pharmacokinetics

Absorption

The absolute oral bioavailability of mefloquine has not been determined since an intravenous formulation is not available. The bioavailability of the tablet formation compared with an oral solution was over 85%. The presence of food significantly enhances the rate and extent of absorption, leading to about

LARIAM (mefloquine hydrochloride)

29 a 40% increase in bioavailability. In healthy volunteers, plasma concentrations
30 peak 6 to 24 hours (median, about 17 hours) after a single dose of Lariam. In a
31 similar group of volunteers, maximum plasma concentrations in µg/L are
32 roughly equivalent to the dose in milligrams (for example, a single 1000 mg
33 dose produces a maximum concentration of about 1000 µg/L). In healthy
34 volunteers, a dose of 250 mg once weekly produces maximum steady-state
35 plasma concentrations of 1000 to 2000 µg/L, which are reached after 7 to 10
36 weeks.

37 Distribution

38 In healthy adults, the apparent volume of distribution is approximately 20
39 L/kg, indicating extensive tissue distribution. Mefloquine may accumulate in
40 parasitized erythrocytes. Experiments conducted in vitro with human blood
41 using concentrations between 50 and 1000 mg/mL showed a relatively
42 constant erythrocyte-to-plasma concentration ratio of about 2 to 1. The
43 equilibrium reached in less than 30 minutes was found to be reversible.
44 Protein binding is about 98%.

45 Mefloquine crosses the placenta. Excretion into breast milk appears to be
46 minimal (see **PRECAUTIONS: Nursing Mothers**).

47 Metabolism

48 Two metabolites have been identified in humans. The main metabolite, 2,8-
49 bis-trifluoromethyl-4-quinoline carboxylic acid, is inactive in *Plasmodium*
50 *falciparum*. In a study in healthy volunteers, the carboxylic acid metabolite
51 appeared in plasma 2 to 4 hours after a single oral dose. Maximum plasma
52 concentrations, which were about 50% higher than those of mefloquine, were
53 reached after 2 weeks. Thereafter, plasma levels of the main metabolite and
54 mefloquine declined at a similar rate. The area under the plasma
55 concentration-time curve (AUC) of the main metabolite was 3 to 5 times
56 larger than that of the parent drug. The other metabolite, an alcohol, was
57 present in minute quantities only.

58 Elimination

59 In several studies in healthy adults, the mean elimination half-life of
60 mefloquine varied between 2 and 4 weeks, with an average of about 3 weeks.
61 Total clearance, which is essentially hepatic, is in the order of 30 mL/min.
62 There is evidence that mefloquine is excreted mainly in the bile and feces. In
63 volunteers, urinary excretion of unchanged mefloquine and its main
64 metabolite under steady-state condition accounted for about 9% and 4% of the
65 dose, respectively. Concentrations of other metabolites could not be measured
66 in the urine.

LARIAM (mefloquine hydrochloride)

67 Pharmacokinetics in Special Clinical Situations

68 *Children and the Elderly*

69 No relevant age-related changes have been observed in the pharmacokinetics
70 of mefloquine. Therefore, the dosage for children has been extrapolated from
71 the recommended adult dose.

72 No pharmacokinetic studies have been performed in patients with renal
73 insufficiency since only a small proportion of the drug is eliminated renally.
74 Mefloquine and its main metabolite are not appreciably removed by
75 hemodialysis. No special chemoprophylactic dosage adjustments are indicated
76 for dialysis patients to achieve concentrations in plasma similar to those in
77 healthy persons.

78 Although clearance of mefloquine may increase in late pregnancy, in general,
79 pregnancy has no clinically relevant effect on the pharmacokinetics of
80 mefloquine.

81 The pharmacokinetics of mefloquine may be altered in acute malaria.

82 Pharmacokinetic differences have been observed between various ethnic
83 populations. In practice, however, these are of minor importance compared
84 with host immune status and sensitivity of the parasite.

85 During long-term prophylaxis (>2 years), the trough concentrations and the
86 elimination half-life of mefloquine were similar to those obtained in the same
87 population after 6 months of drug use, which is when they reached steady
88 state.

89 In vitro and in vivo studies showed no hemolysis associated with glucose-6-
90 phosphate dehydrogenase deficiency (see **ANIMAL TOXICOLOGY**).

91 **Microbiology**

92 Mechanism of Action

93 Mefloquine is an antimalarial agent which acts as a blood schizonticide. Its
94 exact mechanism of action is not known.

95 Activity In Vitro and In Vivo

96 Mefloquine is active against the erythrocytic stages of *Plasmodium* species
97 (see **INDICATIONS AND USAGE**). However, the drug has no effect against
98 the exoerythrocytic (hepatic) stages of the parasite. Mefloquine is effective
99 against malaria parasites resistant to chloroquine (see **INDICATIONS AND**
100 **USAGE**).

101 Drug Resistance

102 Strains of *P. falciparum* with decreased susceptibility to mefloquine can be
103 selected in vitro or in vivo. Resistance of *P. falciparum* to mefloquine has

LARIAM (mefloquine hydrochloride)

104 been reported in areas of multi-drug resistance in South East Asia. Increased
105 incidences of resistance have also been reported in other parts of the world.

106 Cross-Resistance

107 Cross-resistance between mefloquine and halofantrine and cross-resistance
108 between mefloquine and quinine have been observed in some regions.

109 INDICATIONS AND USAGE

110 Treatment of Acute Malaria Infections

111 Lariam is indicated for the treatment of mild to moderate acute malaria caused
112 by mefloquine-susceptible strains of *P. falciparum* (both chloroquine-
113 susceptible and resistant strains) or by *Plasmodium vivax*. There are
114 insufficient clinical data to document the effect of mefloquine in malaria
115 caused by *P. ovale* or *P. malariae*.

116 *Note:* Patients with acute *P. vivax* malaria, treated with Lariam, are at
117 high risk of relapse because Lariam does not eliminate exoerythrocytic
118 (hepatic phase) parasites. To avoid relapse, after initial treatment of the
119 acute infection with Lariam, patients should subsequently be treated
120 with an 8-aminoquinoline derivative (eg, primaquine).

121 Prevention of Malaria

122 Lariam is indicated for the prophylaxis of *P. falciparum* and *P. vivax* malaria
123 infections, including prophylaxis of chloroquine-resistant strains of *P.*
124 *falciparum*.

125 CONTRAINDICATIONS

126 Use of Lariam is contraindicated in patients with a known hypersensitivity to
127 mefloquine or related compounds (eg, quinine and quinidine) or to any of the
128 excipients contained in the formulation. Lariam should not be prescribed for
129 prophylaxis in patients with active depression, a recent history of depression,
130 generalized anxiety disorder, psychosis, or schizophrenia or other major
131 psychiatric disorders, or with a history of convulsions.

132 WARNINGS

133 **In case of life-threatening, serious or overwhelming malaria infections**
134 **due to *P. falciparum*, patients should be treated with an intravenous**
135 **antimalarial drug. Following completion of intravenous treatment,**
136 **Lariam may be given to complete the course of therapy.**

137 **Data on the use of halofantrine subsequent to administration of Lariam**
138 **suggest a significant, potentially fatal prolongation of the QTc interval of**
139 **the ECG. Therefore, halofantrine must not be given simultaneously with**
140 **or subsequent to Lariam. No data are available on the use of Lariam after**
141 **halofantrine (see PRECAUTIONS: Drug Interactions).**

LARIAM (mefloquine hydrochloride)

142 Mefloquine may cause psychiatric symptoms in a number of patients,
143 ranging from anxiety, paranoia, and depression to hallucinations and
144 psychotic behavior. On occasions, these symptoms have been reported to
145 continue long after mefloquine has been stopped. Rare cases of suicidal
146 ideation and suicide have been reported though no relationship to drug
147 administration has been confirmed. To minimize the chances of these
148 adverse events, mefloquine should not be taken for prophylaxis in
149 patients with active depression or with a recent history of depression,
150 generalized anxiety disorder, psychosis, or schizophrenia or other major
151 psychiatric disorders. Lariam should be used with caution in patients
152 with a previous history of depression.

153 During prophylactic use, if psychiatric symptoms such as acute anxiety,
154 depression, restlessness or confusion occur, these may be considered
155 prodromal to a more serious event. In these cases, the drug must be
156 discontinued and an alternative medication should be substituted.

157 Concomitant administration of Lariam and quinine or quinidine may
158 produce electrocardiographic abnormalities.

159 Concomitant administration of Lariam and quinine or chloroquine may
160 increase the risk of convulsions.

161 PRECAUTIONS

162 General

163 Hypersensitivity reactions ranging from mild cutaneous events to anaphylaxis
164 cannot be predicted.

165 In patients with epilepsy, Lariam may increase the risk of convulsions. The
166 drug should therefore be prescribed only for curative treatment in such
167 patients and only if there are compelling medical reasons for its use (see
168 **PRECAUTIONS: Drug Interactions**).

169 Caution should be exercised with regard to activities requiring alertness and
170 fine motor coordination such as driving, piloting aircraft, operating
171 machinery, and deep-sea diving, as dizziness, a loss of balance, or other
172 disorders of the central or peripheral nervous system have been reported
173 during and following the use of Lariam. These effects may occur after therapy
174 is discontinued due to the long half-life of the drug. Lariam should be used
175 with caution in patients with psychiatric disturbances because mefloquine use
176 has been associated with emotional disturbances (see **ADVERSE**
177 **REACTIONS**).

178 In patients with impaired liver function the elimination of mefloquine may be
179 prolonged, leading to higher plasma levels.

180 This drug has been administered for longer than 1 year. If the drug is to be
181 administered for a prolonged period, periodic evaluations including liver

LARIAM (mefloquine hydrochloride)

function tests should be performed. Although retinal abnormalities seen in humans with long-term chloroquine use have not been observed with mefloquine use, long-term feeding of mefloquine to rats resulted in dose-related ocular lesions (retinal degeneration, retinal edema and lenticular opacity at 12.5 mg/kg/day and higher) (see **ANIMAL TOXICOLOGY**). Therefore, periodic ophthalmic examinations are recommended.

Parenteral studies in animals show that mefloquine, a myocardial depressant, possesses 20% of the antifibrillatory action of quinidine and produces 50% of the increase in the PR interval reported with quinine. The effect of mefloquine on the compromised cardiovascular system has not been evaluated. However, transitory and clinically silent ECG alterations have been reported during the use of mefloquine. Alterations included sinus bradycardia, sinus arrhythmia, first degree AV-block, prolongation of the QTc interval and abnormal T waves (see also cardiovascular effects under **PRECAUTIONS: Drug Interactions** and **ADVERSE REACTIONS**). The benefits of Lariam therapy should be weighed against the possibility of adverse effects in patients with cardiac disease.

Laboratory Tests

Periodic evaluation of hepatic function should be performed during prolonged prophylaxis.

Information for Patients

Medication Guide: As required by law, a Lariam Medication Guide is supplied to patients when Lariam is dispensed. An information wallet card is also supplied to patients when Lariam is dispensed. Patients should be instructed to read the Medication Guide when Lariam is received and to carry the information wallet card with them when they are taking Lariam. The complete texts of the Medication Guide and information wallet card are reprinted at the end of this document.

Patients should be advised:

- that malaria can be a life-threatening infection in the traveler;
- that Lariam is being prescribed to help prevent or treat this serious infection;
- that in a small percentage of cases, patients are unable to take this medication because of side effects, and it may be necessary to change medications;
- that when used as prophylaxis, the first dose of Lariam should be taken 1 week prior to arrival in an endemic area;
- that if the patients experience psychiatric symptoms such as acute anxiety, depression, restlessness or confusion, these may be considered prodromal to a more serious event. In these cases, the drug must be discontinued and an alternative medication should be substituted;

LARIAM (mefloquine hydrochloride)

- that no chemoprophylactic regimen is 100% effective, and protective clothing, insect repellents, and bednets are important components of malaria prophylaxis;
- to seek medical attention for any febrile illness that occurs after return from a malarious area and to inform their physician that they may have been exposed to malaria.

Drug Interactions

Drug-drug interactions with Lariam have not been explored in detail. There is one report of cardiopulmonary arrest, with full recovery, in a patient who was taking a beta blocker (propranolol) (see **PRECAUTIONS: General**). The effects of mefloquine on the compromised cardiovascular system have not been evaluated. The benefits of Lariam therapy should be weighed against the possibility of adverse effects in patients with cardiac disease.

Because of the danger of a potentially fatal prolongation of the QTc interval, halofantrine must not be given simultaneously with or subsequent to Lariam (see **WARNINGS**).

Concomitant administration of Lariam and other related compounds (eg, quinine, quinidine and chloroquine) may produce electrocardiographic abnormalities and increase the risk of convulsions (see **WARNINGS**). If these drugs are to be used in the initial treatment of severe malaria, Lariam administration should be delayed at least 12 hours after the last dose. There is evidence that the use of halofantrine after mefloquine causes a significant lengthening of the QTc interval. Clinically significant QTc prolongation has not been found with mefloquine alone.

This appears to be the only clinically relevant interaction of this kind with Lariam, although theoretically, coadministration of other drugs known to alter cardiac conduction (eg, anti-arrhythmic or beta-adrenergic blocking agents, calcium channel blockers, antihistamines or H₁-blocking agents, tricyclic antidepressants and phenothiazines) might also contribute to a prolongation of the QTc interval. There are no data that conclusively establish whether the concomitant administration of mefloquine and the above listed agents has an effect on cardiac function.

In patients taking an anticonvulsant (eg, valproic acid, carbamazepine, phenobarbital or phenytoin), the concomitant use of Lariam may reduce seizure control by lowering the plasma levels of the anticonvulsant. Therefore, patients concurrently taking antiseizure medication and Lariam should have the blood level of their antiseizure medication monitored and the dosage adjusted appropriately (see **PRECAUTIONS: General**).

When Lariam is taken concurrently with oral live typhoid vaccines, attenuation of immunization cannot be excluded. Vaccinations with attenuated live bacteria should therefore be completed at least 3 days before the first dose of Lariam.

LARIAM (mefloquine hydrochloride)

265 No other drug interactions are known. Nevertheless, the effects of Lariam on
266 travelers receiving comedication, particularly diabetics or patients using
267 anticoagulants, should be checked before departure.

268 In clinical trials, the concomitant administration of sulfadoxine and
269 pyrimethamine did not alter the adverse reaction profile.

270 Carcinogenesis, Mutagenesis, Impairment of Fertility

271 Carcinogenesis

272 The carcinogenic potential of mefloquine was studied in rats and mice in 2-
273 year feeding studies at doses of up to 30 mg/kg/day. No treatment-related
274 increases in tumors of any type were noted.

275 Mutagenesis

276 The mutagenic potential of mefloquine was studied in a variety of assay
277 systems including: Ames test, a host-mediated assay in mice, fluctuation tests
278 and a mouse micronucleus assay. Several of these assays were performed with
279 and without prior metabolic activation. In no instance was evidence obtained
280 for the mutagenicity of mefloquine.

281 Impairment of Fertility

282 Fertility studies in rats at doses of 5, 20, and 50 mg/kg/day of mefloquine have
283 demonstrated adverse effects on fertility in the male at the high dose of 50
284 mg/kg/day, and in the female at doses of 20 and 50 mg/kg/day.
285 Histopathological lesions were noted in the epididymides from male rats at
286 doses of 20 and 50 mg/kg/day. Administration of 250 mg/week of mefloquine
287 (base) in adult males for 22 weeks failed to reveal any deleterious effects on
288 human spermatozoa.

289 Pregnancy

290 Teratogenic Effects

291 Pregnancy Category C. Mefloquine has been demonstrated to be teratogenic
292 in rats and mice at a dose of 100 mg/kg/day. In rabbits, a high dose of 160
293 mg/kg/day was embryotoxic and teratogenic, and a dose of 80 mg/kg/day was
294 teratogenic but not embryotoxic. There are no adequate and well-controlled
295 studies in pregnant women. However, clinical experience with Lariam has not
296 revealed an embryotoxic or teratogenic effect. Mefloquine should be used
297 during pregnancy only if the potential benefit justifies the potential risk to the
298 fetus. Women of childbearing potential who are traveling to areas where
299 malaria is endemic should be warned against becoming pregnant. Women of
300 childbearing potential should also be advised to practice contraception during
301 malaria prophylaxis with Lariam and for up to 3 months thereafter. However,
302 in the case of unplanned pregnancy, malaria chemoprophylaxis with Lariam is
303 not considered an indication for pregnancy termination.

LARIAM (mefloquine hydrochloride)

304 Nursing Mothers

305 Mefloquine is excreted in human milk in small amounts, the activity of which
306 is unknown. Based on a study in a few subjects, low concentrations (3% to
307 4%) of mefloquine were excreted in human milk following a dose equivalent
308 to 250 mg of the free base. Because of the potential for serious adverse
309 reactions in nursing infants from mefloquine, a decision should be made
310 whether to discontinue the drug, taking into account the importance of the
311 drug to the mother.

312 Pediatric Use

313 Use of Lariam to treat acute, uncomplicated *P. falciparum* malaria in pediatric
314 patients is supported by evidence from adequate and well-controlled studies of
315 Lariam in adults with additional data from published open-label and
316 comparative trials using Lariam to treat malaria caused by *P. falciparum* in
317 patients younger than 16 years of age. The safety and effectiveness of Lariam
318 for the treatment of malaria in pediatric patients below the age of 6 months
319 have not been established.

320 In several studies, the administration of Lariam for the treatment of malaria
321 was associated with early vomiting in pediatric patients. Early vomiting was
322 cited in some reports as a possible cause of treatment failure. If a second dose
323 is not tolerated, the patient should be monitored closely and alternative
324 malaria treatment considered if improvement is not observed within a
325 reasonable period of time (see **DOSAGE AND ADMINISTRATION**).

326 Geriatric Use

327 Clinical studies of Lariam did not include sufficient numbers of subjects aged
328 65 and over to determine whether they respond differently from younger
329 subjects. Other reported clinical experience has not identified differences in
330 responses between the elderly and younger patients. Since
331 electrocardiographic abnormalities have been observed in individuals treated
332 with Lariam (see **PRECAUTIONS**) and underlying cardiac disease is more
333 prevalent in elderly than in younger patients, the benefits of Lariam therapy
334 should be weighed against the possibility of adverse cardiac effects in elderly
335 patients.

336 ADVERSE REACTIONS

337 Clinical

338 At the doses used for treatment of acute malaria infections, the symptoms
339 possibly attributable to drug administration cannot be distinguished from
340 those symptoms usually attributable to the disease itself.

341 Among subjects who received mefloquine for prophylaxis of malaria, the
342 most frequently observed adverse experience was vomiting (3%). Dizziness,

LARIAM (mefloquine hydrochloride)

343 syncope, extrasystoles and other complaints affecting less than 1% were also
344 reported.

345 Among subjects who received mefloquine for treatment, the most frequently
346 observed adverse experiences included: dizziness, myalgia, nausea, fever,
347 headache, vomiting, chills, diarrhea, skin rash, abdominal pain, fatigue, loss of
348 appetite, and tinnitus. Those side effects occurring in less than 1% included
349 bradycardia, hair loss, emotional problems, pruritus, asthenia, transient
350 emotional disturbances and telogen effluvium (loss of resting hair). Seizures
351 have also been reported.

352 Two serious adverse reactions were cardiopulmonary arrest in one patient
353 shortly after ingesting a single prophylactic dose of mefloquine while
354 concomitantly using propranolol (see **PRECAUTIONS: Drug Interactions**),
355 and encephalopathy of unknown etiology during prophylactic mefloquine
356 administration. The relationship of encephalopathy to drug administration
357 could not be clearly established.

358 Postmarketing

359 Postmarketing surveillance indicates that the same kind of adverse
360 experiences are reported during prophylaxis, as well as acute treatment.

361 The most frequently reported adverse events are nausea, vomiting, loose
362 stools or diarrhea, abdominal pain, dizziness or vertigo, loss of balance, and
363 neuropsychiatric events such as headache, somnolence, and sleep disorders
364 (insomnia, abnormal dreams). These are usually mild and may decrease
365 despite continued use.

366 Occasionally, more severe neuropsychiatric disorders have been reported such
367 as: sensory and motor neuropathies (including paresthesia, tremor and ataxia),
368 convulsions, agitation or restlessness, anxiety, depression, mood changes,
369 panic attacks, forgetfulness, confusion, hallucinations, aggression, psychotic
370 or paranoid reactions and encephalopathy. Rare cases of suicidal ideation and
371 suicide have been reported though no relationship to drug administration has
372 been confirmed.

373 Other infrequent adverse events include:

374 *Cardiovascular Disorders:* circulatory disturbances (hypotension,
375 hypertension, flushing, syncope), chest pain, tachycardia or palpitation,
376 bradycardia, irregular pulse, extrasystoles, A-V block, and other transient
377 cardiac conduction alterations

378 *Skin Disorders:* rash, exanthema, erythema, urticaria, pruritus, edema, hair
379 loss, erythema multiforme, and Stevens-Johnson syndrome

380 *Musculoskeletal Disorders:* muscle weakness, muscle cramps, myalgia, and
381 arthralgia

LARIAM (mefloquine hydrochloride)

382 *Respiratory Disorders:* dyspnea, pneumonitis of possible allergic etiology

383 *Other Symptoms:* visual disturbances, vestibular disorders including tinnitus
384 and hearing impairment, asthenia, malaise, fatigue, fever, sweating, chills,
385 dyspepsia and loss of appetite

Laboratory

387 The most frequently observed laboratory alterations which could be possibly
388 attributable to drug administration were decreased hematocrit, transient
389 elevation of transaminases, leukopenia and thrombocytopenia. These
390 alterations were observed in patients with acute malaria who received
391 treatment doses of the drug and were attributed to the disease itself.

392 During prophylactic administration of mefloquine to indigenous populations
393 in malaria-endemic areas, the following occasional alterations in laboratory
394 values were observed: transient elevation of transaminases, leukocytosis or
395 thrombocytopenia.

396 Because of the long half-life of mefloquine, adverse reactions to Lariam may
397 occur or persist up to several weeks after the last dose.

OVERDOSAGE

Symptoms and Signs

400 In cases of overdosage with Lariam, the symptoms mentioned under
401 **ADVERSE REACTIONS** may be more pronounced.

Treatment

403 Patients should be managed by symptomatic and supportive care following
404 Lariam overdose. There are no specific antidotes. Monitor cardiac function (if
405 possible by ECG) and neuropsychiatric status for at least 24 hours. Provide
406 symptomatic and intensive supportive treatment as required, particularly for
407 cardiovascular disturbances.

DOSAGE AND ADMINISTRATION (see INDICATIONS AND USAGE)

Adult Patients

410 Treatment of mild to moderate malaria in adults caused by *P. vivax* or
411 mefloquine-susceptible strains of *P. falciparum*

412 Five tablets (1250 mg) mefloquine hydrochloride to be given as a single oral
413 dose. The drug should not be taken on an empty stomach and should be
414 administered with at least 8 oz (240 mL) of water.

415 If a full-treatment course with Lariam does not lead to improvement within 48
416 to 72 hours, Lariam should not be used for retreatment. An alternative therapy
417 should be used. Similarly, if previous prophylaxis with mefloquine has failed,
418 Lariam should not be used for curative treatment.

LARIAM (mefloquine hydrochloride)

419 *Note:* Patients with acute *P. vivax* malaria, treated with Lariam, are at
420 high risk of relapse because Lariam does not eliminate exoerythrocytic
421 (hepatic phase) parasites. To avoid relapse after initial treatment of the
422 acute infection with Lariam, patients should subsequently be treated
423 with an 8-aminoquinoline derivative (eg, primaquine).

424 Malaria Prophylaxis

425 One 250 mg Lariam tablet once weekly.

426 Prophylactic drug administration should begin 1 week before arrival in an
427 endemic area. Subsequent weekly doses should be taken regularly, always on
428 the same day of each week, preferably after the main meal. To reduce the risk
429 of malaria after leaving an endemic area, prophylaxis must be continued for 4
430 additional weeks to ensure suppressive blood levels of the drug when
431 merozoites emerge from the liver. Tablets should not be taken on an empty
432 stomach and should be administered with at least 8 oz (240 mL) of water.

433 In certain cases, eg, when a traveler is taking other medication, it may be
434 desirable to start prophylaxis 2 to 3 weeks prior to departure, in order to
435 ensure that the combination of drugs is well tolerated (see **PRECAUTIONS:**
436 **Drug Interactions**).

437 When prophylaxis with Lariam fails, physicians should carefully evaluate
438 which antimalarial to use for therapy.

439 Pediatric Patients

440 Treatment of mild to moderate malaria in pediatric patients caused by
441 mefloquine-susceptible strains of *P. falciparum*

442 Twenty (20) to 25 mg/kg body weight. Splitting the total therapeutic dose into
443 2 doses taken 6 to 8 hours apart may reduce the occurrence or severity of
444 adverse effects. Experience with Lariam in infants less than 3 months old or
445 weighing less than 5 kg is limited. The drug should not be taken on an empty
446 stomach and should be administered with ample water. The tablets may be
447 crushed and suspended in a small amount of water, milk or other beverage for
448 administration to small children and other persons unable to swallow them
449 whole.

450 If a full-treatment course with Lariam does not lead to improvement within 48
451 to 72 hours, Lariam should not be used for retreatment. An alternative therapy
452 should be used. Similarly, if previous prophylaxis with mefloquine has failed,
453 Lariam should not be used for curative treatment.

454 In pediatric patients, the administration of Lariam for the treatment of malaria
455 has been associated with early vomiting. In some cases, early vomiting has
456 been cited as a possible cause of treatment failure (see **PRECAUTIONS**). If a
457 significant loss of drug product is observed or suspected because of vomiting,
458 a second full dose of Lariam should be administered to patients who vomit

LARIAM (mefloquine hydrochloride)

less than 30 minutes after receiving the drug. If vomiting occurs 30 to 60 minutes after a dose, an additional half-dose should be given. If vomiting recurs, the patient should be monitored closely and alternative malaria treatment considered if improvement is not observed within a reasonable period of time.

The safety and effectiveness of Lariam to treat malaria in pediatric patients below the age of 6 months have not been established.

Malaria Prophylaxis

The following doses have been extrapolated from the recommended adult dose. Neither the pharmacokinetics, nor the clinical efficacy of these doses have been determined in children owing to the difficulty of acquiring this information in pediatric subjects. The recommended prophylactic dose of Lariam is approximately 5 mg/kg body weight once weekly. One 250 mg Lariam tablet should be taken once weekly in pediatric patients weighing over 45 kg. In pediatric patients weighing less than 45 kg, the weekly dose decreases in proportion to body weight:

30 to 45 kg: 3/4 tablet

20 to 30 kg: 1/2 tablet

10 to 20 kg: 1/4 tablet

5 to 10 kg: 1/8 tablet*

*Approximate tablet fraction based on a dosage of 5 mg/kg body weight. Exact doses for children weighing less than 10 kg may best be prepared and dispensed by pharmacists.

Experience with Lariam in infants less than 3 months old or weighing less than 5 kg is limited.

HOW SUPPLIED

Lariam is available as scored, white, round tablets, containing 250 mg of mefloquine hydrochloride in unit-dose packages of 25 (NDC 0004-0172-02). Imprint on tablets: LARIAM 250 ROCHE

Tablets should be stored at 25°C (77°F); excursions permitted to 15° to 30°C (59° to 86°F).

ANIMAL TOXICOLOGY

Ocular lesions were observed in rats fed mefloquine daily for 2 years. All surviving rats given 30 mg/kg/day had ocular lesions in both eyes characterized by retinal degeneration, opacity of the lens, and retinal edema. Similar but less severe lesions were observed in 80% of female and 22% of male rats fed 12.5 mg/kg/day for 2 years. At doses of 5 mg/kg/day, only corneal lesions were observed. They occurred in 9% of rats studied.

LARIAM (mefloquine hydrochloride)

497 Revised: May 2008

498 MEDICATION GUIDE

499 **This Medication Guide is intended only for travelers who are taking**
500 **Lariam to prevent malaria.** The information may not apply to patients who
501 are sick with malaria and who are taking Lariam to treat malaria.

502 An information wallet card is provided with this Medication Guide. Carry it
503 with you when you are taking Lariam.

504 This Medication Guide was revised in May 2004. Please read it before you
505 start taking Lariam and each time you get a refill. There may be new
506 information. This Medication Guide does not take the place of talking with
507 your prescriber (doctor or other health care provider) about Lariam and
508 malaria prevention. Only you and your prescriber can decide if Lariam is right
509 for you. If you cannot take Lariam, you may be able to take a different
510 medicine to prevent malaria.

511 What is the most important information I should know about 512 **Lariam?**

513 1. Take Lariam exactly as prescribed to prevent malaria.

514 Malaria is an infection that can cause death and is spread to humans
515 through mosquito bites. If you travel to parts of the world where the
516 mosquitoes carry the malaria parasite, you must take a malaria prevention
517 medicine. Lariam is one of a small number of medications approved to
518 prevent and to treat malaria. If taken correctly, Lariam is effective at
519 preventing malaria but, like all medications, it may produce side effects in
520 some patients.

521 2. Lariam can rarely cause serious mental problems in some patients.

522 The most frequently reported side effects with Lariam, such as nausea,
523 difficulty sleeping, and bad dreams are usually mild and do not cause
524 people to stop taking the medicine. However, people taking Lariam
525 occasionally experience severe anxiety, feelings that people are against
526 them, hallucinations (seeing or hearing things that are not there, for
527 example), depression, unusual behavior, or feeling disoriented. There have
528 been reports that in some patients these side effects continue after Lariam
529 is stopped. Some patients taking Lariam think about killing themselves,
530 and there have been rare reports of suicides. It is not known whether
531 Lariam was responsible for these suicides.

532 3. You need to take malaria prevention medicine before you travel to a 533 **malaria area, while you are in a malaria area, and after you return** 534 **from a malaria area.**

535 Medicines approved in the United States for malaria prevention include
536 Lariam, doxycycline, atovaquone/proguanil, hydroxychloroquine, and

LARIAM (mefloquine hydrochloride)

537 chloroquine. Not all of these drugs work equally as well in all areas of the
538 world where there is malaria. The chloroquines, for example, do not work
539 in areas where the malaria parasite has developed resistance to
540 chloroquine. Lariam may be effective against malaria that is resistant to
541 chloroquine or other drugs. All drugs to treat malaria have side effects that
542 are different for each one. For example, some may make your skin more
543 sensitive to sunlight (Lariam does not do this). However, if you use
544 Lariam to prevent malaria and you develop a sudden onset of anxiety,
545 depression, restlessness, confusion (possible signs of more serious mental
546 problems), or you develop other serious side effects, contact a doctor or
547 other health care provider. It may be necessary to stop taking Lariam and
548 use another malaria prevention medicine instead. If you can't get another
549 medicine, leave the malaria area. However, be aware that leaving the
550 malaria area may not protect you from getting malaria. You still need to
551 take a malaria prevention medicine.

552 Who should not take Lariam?

553 Do not take Lariam to **prevent** malaria if you

- 554 • **have depression or had depression recently**
- 555 • **have had recent mental illness or problems**, including anxiety disorder,
556 schizophrenia (a severe type of mental illness), or psychosis (losing touch
557 with reality)
- 558 • **have or had seizures (epilepsy or convulsions)**
- 559 • **are allergic to quinine or quinidine (medicines related to Lariam)**

560 **Tell your prescriber about all your medical conditions.** Lariam may not be
561 right for you if you have certain conditions, especially the ones listed below:

- 562 • **Heart disease.** Lariam may not be right for you.
- 563 • **Pregnancy.** Tell your prescriber if you are pregnant or plan to become
564 pregnant. It is dangerous for the mother and for the unborn baby (fetus) to
565 get malaria during pregnancy. Therefore, ask your prescriber if you should
566 take Lariam or another medicine to prevent malaria while you are
567 pregnant.
- 568 • **Breast-feeding.** Lariam can pass through your milk and may harm the
569 baby. Therefore, ask your prescriber whether you will need to stop breast-
570 feeding or use another medicine.
- 571 • **Liver problems.**

572 **Tell your prescriber about all the medicines you take, including**
573 **prescription and non-prescription medicines, vitamins, and herbal**

LARIAM (mefloquine hydrochloride)

574 **supplements.** Some medicines may give you a higher chance of having
575 serious side effects from Lariam.

576 How should I take Lariam?

577 **Take Lariam exactly as prescribed. If you are an adult or pediatric**
578 **patient weighing 45 kg (99 pounds) or less, your prescriber will tell you**
579 **the correct dose based on your weight.**

580 To prevent malaria

- 581 • For adults and pediatric patients weighing over 45 kg, take 1 tablet of
582 Lariam at least 1 week before you travel to a malaria area (or 2 to 3 weeks
583 before you travel to a malaria area, if instructed by your prescriber). This
584 starts the prevention and also helps you see how Lariam affects you and
585 the other medicines you take. **Take 1 Lariam tablet once a week,** on the
586 same day each week, while in a malaria area.
- 587 • **Continue taking Lariam for 4 weeks after returning from a malaria**
588 **area.** If you cannot continue taking Lariam due to side effects or for other
589 reasons, contact your prescriber.
- 590 • Take Lariam just after a meal and with at least 1 cup (8 ounces) of water.
- 591 • For children, Lariam can be given with water or crushed and mixed with
592 water or sugar water. The prescriber will tell you the correct dose for
593 children based on the child's weight.
- 594 • If you are told by a doctor or other health care provider to stop taking
595 Lariam due to side effects or for other reasons, it will be necessary to take
596 another malaria medicine. You must take **malaria prevention medicine**
597 **before you travel to a malaria area, while you are in a malaria area,**
598 **and after you return from a malaria area. If you don't have access to**
599 **a doctor or other health care provider or to another medicine besides**
600 **Lariam and have to stop taking it, leave the malaria area. However, be**
601 **aware that leaving the malaria area may not protect you from getting**
602 **malaria. You still need to take a malaria prevention medicine.**

603 What should I avoid while taking Lariam?

- 604 • **Halofantrine (marketed under various brand names),** a medicine used
605 to treat malaria. Taking both of these medicines together can cause serious
606 heart problems that can cause death.
- 607 • **Do not become pregnant.** Women should use effective birth control
608 while taking Lariam.
- 609 • **Quinine, quinidine, or chloroquine (other medicines used to treat**
610 **malaria).** Taking these medicines with Lariam could cause changes in
611 your heart rate or increase the risk of seizures.

LARIAM (mefloquine hydrochloride)

In addition:

- **Be careful driving or in other activities** needing alertness and careful movements (fine motor coordination). Lariam can cause dizziness or loss of balance, even after you stop taking it.
- **Be aware that certain vaccines may not work if given while you are taking Lariam.** Your prescriber may want you to finish taking your vaccines at least 3 days before starting Lariam.

What are the possible side effects of Lariam?

Lariam, like all medicines, may cause side effects in some patients. The most frequently reported side effects with Lariam when used for prevention of malaria include nausea, vomiting, diarrhea, dizziness, difficulty sleeping, and bad dreams. These are usually mild and do not cause people to stop taking the medicine.

Lariam may cause serious mental problems in some patients (see “What is the most important information I should know about Lariam?”).

Lariam may affect your liver and your eyes if you take it for a long time. Your prescriber will tell you if you should have your eyes and liver checked while taking Lariam.

What else should I know about preventing malaria?

- **Find out whether you need malaria prevention.** Before you travel, talk with your prescriber about your travel plans to determine whether you need to take medicine to prevent malaria. Even in those countries where malaria is present, there may be areas of the country that are free of malaria. In general, malaria is more common in rural (country) areas than in big cities, and it is more common during rainy seasons, when mosquitoes are most common. You can get information about the areas of the world where malaria occurs from the Centers for Disease Control and Prevention (CDC) and from local authorities in the countries you visit. If possible, plan your travel to reduce the risk of malaria.
- **Take medicine to prevent malaria infection.** Without malaria prevention medicine, you have a higher risk of getting malaria. Malaria starts with flu-like symptoms, such as chills, fever, muscle pains, and headaches. However, malaria can make you very sick or cause death if you don’t seek medical help immediately. These symptoms may disappear for a while, and you may think you are well. But, the symptoms return later and then it may be too late for successful treatment.

Malaria can cause confusion, coma, and seizures. It can cause kidney failure, breathing problems, and severe damage to red blood cells. However, malaria can be easily diagnosed with a blood test, and if caught in time, can be effectively treated.

LARIAM (mefloquine hydrochloride)

652 **If you get flu-like symptoms (chills, fever, muscle pains, or**
653 **headaches) after you return from a malaria area,** get medical help
654 right away and tell your prescriber that you may have been exposed to
655 malaria.

656 People who have lived for many years in areas with malaria may have
657 some immunity to malaria (they do not get it as easily) and may not
658 take malaria prevention medicine. This does not mean that you don't
659 need to take malaria prevention medicine.

660 • **Protect against mosquito bites.** Medicines do not always completely
661 prevent your catching malaria from mosquito bites. So protect yourself
662 very well against mosquitoes. Cover your skin with long sleeves and long
663 pants, and use mosquito repellent and bednets while in malaria areas. If
664 you are out in the bush, you may want to pre-wash your clothes with
665 permethrin. This is a mosquito repellent that may be effective for weeks
666 after use. Ask your prescriber for other ways to protect yourself.

667 **General information about the safe and effective use of Lariam.**

668 Medicines are sometimes prescribed for conditions not listed in Medication
669 Guides. If you have any concerns about Lariam, ask your prescriber. This
670 Medication Guide contains certain important information for travelers visiting
671 areas with malaria. Your prescriber or pharmacist can give you information
672 about Lariam that was written for health care professionals. Do not use
673 Lariam for a condition for which it was not prescribed. Do not share Lariam
674 with other people.

675 This Medication Guide has been approved by the U.S. Food and Drug
676 Administration.

677 Medication Guide Revised: May 2004

678

679 Reprint of information wallet card:

LARIAM (mefloquine hydrochloride)

680

 Lariam (mefloquine hydrochloride) Tablets Carry this information wallet card with you when you are taking Lariam.	
<p>You need to take malaria prevention medicine before you travel to a malaria area, while you are in a malaria area, and after you return from a malaria area.</p> <p>If taken correctly, Lariam is effective at preventing malaria but, like all medications, it may produce side effects in some patients.</p> <p>If you use Lariam to prevent malaria and you develop a sudden onset of anxiety, depression, restlessness, confusion (possible signs of more serious mental problems), or you develop other serious side effects, contact a doctor or other health care provider. It may be necessary to stop taking Lariam and use another malaria prevention medicine instead.</p>	<p>Other medicines approved in the United States for malaria prevention include: doxycycline, atovaquone/proguanil, hydroxychloroquine, and chloroquine. Not all malaria medicines work equally well in malaria areas. The chloroquines, for example, do not work in many parts of the world. If you can't get another medicine, leave the malaria area. However, be aware that leaving the malaria area may not protect you from getting malaria. You still need to take a malaria prevention medicine.</p> <p>Please read the Medication Guide for additional information on Lariam.</p> <p style="text-align: right;">Card Revised: May 2004</p>

681

682

683

684

685

686

Manufactured by:
F. HOFFMANN-LA ROCHE LTD
Basel, Switzerland

Distributed by:



Pharmaceuticals

687

Roche Laboratories Inc.
340 Kingsland Street
Nutley, New Jersey 07110-1199

688

Copyright © 1999-2008 by Roche Laboratories Inc. All rights reserved.